

Low Variance at large scales of WMAP 9 year data

A. Gruppuso,^{1,2,*} P. Natoli,^{3,4,1} F. Paci,⁵ F. Finelli,^{1,2}

D. Molinari,⁶ A. De Rosa,¹ and N. Mandolesi^{1,3,7}

¹*INAF-IASF Bologna, Via P. Gobetti 101, I-40129, Bologna, Italy*

²*INFN, Sezione di Bologna, Via Irnerio 46, I-40126 Bologna, Italy*

³*Dipartimento di Fisica e Scienze della Terra,*

Università degli Studi di Ferrara e Sezione INFN di Ferrara, via Saragat 1, I-44100 Ferrara, Italy

⁴*Agenzia Spaziale Italiana Science Data Center,*

c/o ESRIN, via Galileo Galilei, Frascati, Italy

⁵*SISSA - Scuola Internazionale Superiore di Studi Avanzati,*

via Bonomea 265, I-34136 Trieste, Italy

⁶*Dipartimento di Astronomia, Università degli Studi di Bologna,*

Via Camillo Ranzani 1, I-40127 Bologna, Italy

⁷*Agenzia Spaziale Italiana, Viale del Politecnico, I-00198 Roma, Italy*

We use an optimal estimator to study the variance of the WMAP 9 CMB field at low resolution, in both temperature and polarization. Employing realistic Monte Carlo simulation, we find statistically significant deviations from the Λ CDM model in several sky cuts for the temperature field. For the largest mask considered in this analysis, which leaves 46% of the sky, the CMB sky and Λ CDM are incompatible at $> 99.94\%$ C.L. at large angles ($> 5^\circ$). We find instead no anomaly in polarization. As a byproduct of our analysis, we present new, optimal estimates of the WMAP 9 CMB angular power spectra from the WMAP 9 year data at low resolution.

I. INTRODUCTION

The cosmic microwave background (CMB) anisotropy field can be used to probe cosmology to high precision, as shown by the Wilkinson Microwave Anisotropy Probe (WMAP) 9 years results [13] and by the very recent PLANCK cosmological results (see [21] and references therein). CMB data have given a significant contribution in setting up the Λ cold dark matter (Λ CDM) cosmological concordance model. The latter establishes a set of basic quantities for which CMB observations and other cosmological and astrophysical data-sets agree¹: spatial curvature close to zero; $\sim 68.5\%$ of the cosmic density in the form of Dark Energy; $\sim 26.5\%$ in cold dark matter;

* gruppuso@iasfbo.inaf.it

¹ See, however, [24] for a tension concerning the Ω_m estimate from PLANCK CMB and galaxy clusters data.

$\sim 5\%$ in baryonic matter; and non perfectly scale invariant adiabatic, primordial perturbations compatible with Gaussianity [23]. The largest angular scales of the CMB anisotropies have remained largely primordial, having evolved only in amplitude since the end of the inflation. They can be used to probe the physics of the very early Universe. Accurate estimates of their amplitude is a crucial to extract information about such primordial phases [25].

In this paper we provide new, temperature and polarization, estimates of the CMB Angular Power Spectra (APS) from the WMAP 9 year low resolution data using a Quadratic Maximum Likelihood (QML) method [31]. We have used *BolPol*, an efficient, fully polarized, implementation of the QML method. *BolPol* has been already applied to WMAP 5 [10] and WMAP 7 [11] low resolution data. At large angular scales, QML estimates are superior to the pseudo spectrum techniques normally used in the CMB world [9, 14, 27, 28] because of their smaller intrinsic variance [17]. The WMAP 9 analysis has been derived using a suboptimal implementation of the QML method that extends to high multipole [1]. *BolPol* is fully optimal, but computational considerations limit its application to smaller multipoles. This is fully adequate for the low resolution analysis presented here. For other applications of our QML implementation see [19, 20], where it is used to constrain the CMB hemispherical power asymmetry in WMAP 5 and 7 data respectively, and [12], where the low resolution spectrum of the cosmological birefringence angle has been derived for the first time².

In this paper, we provide new APS estimates from WMAP 9 year low resolution data. At the same time, we check for their stability against Galactic masking. Specifically, we look at the variance of the temperature and polarization maps. The variance of the CMB temperature field has been studied in previous works (see [3] for WMAP 5 and 7 year data and [18] for WMAP 3 year data) which, however, focused on the small scale pattern. Here we focus on the lowest multipoles of WMAP 9 year data, we adopt a new estimator (based on *BolPol*) and consider also polarization.

The paper is organized as follows. In Section II we briefly describe the QML method and our implementation, *BolPol*. Section III is devoted to the description of the WMAP 9 year low resolution data set under consideration. APS estimates of this data set are given in Section IV where the consistency of all the spectra with what expected in the Λ CDM model is evaluated through a χ^2 analysis. In the same Section the stability of the obtained estimates is studied against the Galactic masking. The analysis of the variance is presented in Section V. Our conclusions are drawn in Section VI where some speculations about a possible origin for this

² See also [29], where *BolISW*, a code largely based on *BolPol*, has been used to detect the ISW effect

effect are set forth.

II. ANGULAR POWER SPECTRA ESTIMATOR

In order to evaluate the APS we adopt the QML estimator, introduced in [30] and extended to polarization in [31]. In this section we describe the essence of such a method.

Given a map in temperature and polarization $\mathbf{x} = (\mathbf{T}, \mathbf{Q}, \mathbf{U})$, the QML provides estimates $\hat{C}_\ell^X$ - with X being one of TT, EE, TE, BB, TB, EB - of the APS as:

$$\hat{C}_\ell^X = \sum_{\ell', X'} (F^{-1})_{\ell\ell'}^{XX'} \left[\mathbf{x}^t \mathbf{E}_{X'}^{\ell\ell'} \mathbf{x} - \text{tr}(\mathbf{N} \mathbf{E}_{X'}^{\ell\ell'}) \right], \quad (1)$$

where the $F_{XX'}^{\ell\ell'}$ is the Fisher matrix, defined as

$$F_{XX'}^{\ell\ell'} = \frac{1}{2} \text{tr} \left[\mathbf{C}^{-1} \frac{\partial \mathbf{C}}{\partial C_\ell^X} \mathbf{C}^{-1} \frac{\partial \mathbf{C}}{\partial C_{\ell'}^{X'}} \right], \quad (2)$$

and the $\mathbf{E}_X^\ell$ matrix is given by

$$\mathbf{E}_X^\ell = \frac{1}{2} \mathbf{C}^{-1} \frac{\partial \mathbf{C}}{\partial C_\ell^X} \mathbf{C}^{-1}, \quad (3)$$

with $\mathbf{C} = \mathbf{S}(C_\ell^X) + \mathbf{N}$ being the global covariance matrix (signal plus noise contribution).

Although an initial assumption for a fiducial power spectrum C_ℓ^X is needed, the QML method provides unbiased estimates of the power spectrum contained in the map regardless of the initial guess,

$$\langle \hat{C}_\ell^X \rangle = \bar{C}_\ell^X, \quad (4)$$

where the average is taken over the ensemble of realizations (or, in a practical test, over Monte Carlo realizations extracted from $\bar{C}_\ell^X$). On the other hand, the covariance matrix associated to the estimates,

$$\langle \Delta \hat{C}_\ell^X \Delta \hat{C}_{\ell'}^{X'} \rangle = (F^{-1})_{\ell\ell'}^{XX'}, \quad (5)$$

does depend on the initial assumption for C_ℓ^X : the closer the guess to the true power spectrum is, the closer are the error bars to minimum variance. According to the Cramer-Rao inequality, which sets a limit to the accuracy of an estimator, Eq. (5) tells us that the QML has the smallest error bars. The QML is then an ‘optimal’ estimator.

This method has been implemented in a F90 code, named *BolPol*. Further details can be found in [10].

III. DATA SET

In this Section we describe the WMAP 9 year data set that we have considered. We use the temperature ILC map smoothed at 9.1285 degrees and reconstructed at HealPix³ [8] resolution $N_{side} = 16$, the foreground cleaned low resolution maps and the noise covariance matrix in (Q, U) publicly available at the LAMBDA website⁴ for the frequency channels Ka, Q and W as considered by [1] and [16] for the low ℓ analysis. These frequency channels have been co-added as follows [11, 15]

$$m_{tot} = C_{tot}(C_{Ka}^{-1}m_{Ka} + C_Q^{-1}m_Q + C_V^{-1}m_V), \quad (6)$$

where m_i , C_i are the polarization maps and covariances (for $i = Ka, Q$ and V) and

$$C_{tot}^{-1} = C_{Ka}^{-1} + C_Q^{-1} + C_V^{-1}. \quad (7)$$

This polarization data set has been extended to temperature considering the ILC map. We have added to the temperature map a random noise realization with variance of $1\mu K^2$ as suggested in [7]. Consistently, the noise covariance matrix for TT is taken to be diagonal with variance equal to $1\mu K^2$.

IV. ESTIMATES

Taking into account the kq85 mask for the temperature map⁵ and the P06 mask for polarization maps we estimate all the 6 CMB spectra of WMAP 9 year data with *BolPol*, see red symbols of Fig. 1. In the same Figure we also show the averages of 10000 “CMB plus noise” Monte-Carlo (MC) simulations (see black symbols) where the CMB is randomly extracted from a Λ CDM model, drawn as dotted line, and the noise is obtained through a Cholesky decomposition of the noise covariance matrix. Consistently with WMAP data, also the temperature simulated maps have been smoothed at that scale, i.e. 9.1285 degrees.

A. χ^2 consistency with Λ CDM model

In order to evaluate the consistency with the Λ CDM model (identified in the following with a set of C_ℓ^{th} for each spectrum) we adopt a frequentist approach, see Fig. 2. For each spectrum

³ <http://healpix.jpl.nasa.gov/>

⁴ <http://lambda.gsfc.nasa.gov/>

⁵ Best-fit monopole and dipole have been subtracted from the observed ILC map through the HealPix routine REMOVE-DIPOLE [8].




FIG. 1. WMAP 9 year CMB spectra at low resolution. ILC in Temperature, masked with kq85 mask and the co-addition of Ka, Q and V channel in Polarization masked with the P06 mask. From top to bottom, TT, EE, TE, BB, TB and EB spectrum. For each spectrum we show $\ell(\ell+1)C_\ell/2\pi$ (given in μK^2) versus the multipole order ℓ . Red estimates are for the WMAP 9 year maps. Black estimates are the averages of 10000 Monte-Carlo simulations where the CMB is randomly extracted from a Λ CDM model (shown as dotted lines) and the noise is obtained through a Cholesky decomposition of the noise covariance matrix. See also the text.




FIG. 2. χ^2 test for consistency with Λ CDM model. Each panel shows the counts versus χ^2 . The histograms are built through Monte Carlo simulations. The red lines are the χ^2 values we obtain for the WMAP 9 year low resolution data. See also the text.

(TT, EE, TE, BB, TB and EB) we plot the histogram of the χ_i^2 values defined as

$$\chi_i^2 = \sum_{\ell, \ell'=2,32} (C_{\ell,i} - C_{\ell}^{th}) M_{\ell\ell'}^{-1} (C_{\ell',i} - C_{\ell'}^{th}), \quad (8)$$

where $i = 1, 10000$ runs over the MC realizations and where the covariance matrix $M_{\ell\ell'}$ is defined as

$$M_{\ell\ell'} = \langle (C_{\ell,i} - C_{\ell}^{th})(C_{\ell',i} - C_{\ell'}^{th}) \rangle, \quad (9)$$

with $\langle \cdot \cdot \rangle$ standing for the ensemble average. These histograms will be compared with the χ_{WMAP9}^2 obtained with the WMAP 9 year data for each spectrum, defined as

$$\chi_{\text{WMAP9}}^2 = \sum_{\ell, \ell'=2,32} (C_{\ell,\text{W9}} - C_{\ell}^{th}) M_{\ell\ell'}^{-1} (C_{\ell',\text{W9}} - C_{\ell'}^{th}), \quad (10)$$

where $C_{\ell,\text{W9}}$ are the WMAP 9 year spectra. Fig. 2 shows that the WMAP 9 year low resolution data appear to be well consistent with the Λ CDM model defined by the WMAP 9 fiducial model, represented by the dotted line shown in Fig. 1. See Table I for a quantitative analysis.

TABLE I. Percentage to obtain a χ^2 value from a random extraction of Λ CDM model smaller than what observed by WMAP 9.

Spectrum Percentage	
TT	59.26
EE	88.99
TE	59.25
BB	46.13
TB	12.22
EB	85.73

TABLE II. Sky fraction of the sky coverage of the considered masks. See also Fig. 3.

Case	Extension wrt kq85 and P06	Sky fraction in T	Sky fraction in P
a	+0 deg	0.78	0.73
b	+4 deg	0.68	0.65
c	+8 deg	0.56	0.53
d	+12 deg	0.46	0.43

B. Stability of the results versus Galactic masking

In this Subsection we test the stability of the WMAP 9 year data estimates considering various Galactic masks, see Fig. 3. More specifically, we extend the edges of the kq85 temperature mask and the P06 polarization mask by 4, 8 and 12 degrees. In Table II we provide the sky fraction for each considered case and for reference we rename with the letters “a”, “b”, “c” and “d” the four cases we take into account. Case “a” is what is already considered in Subsection IV A.

In Fig. 4 we show the six spectra for all the four considered cases. The overall consistency is good. However, the lowest multipoles of the TT spectrum show a clear trend: increasing the temperature mask, we obtain a lowering of the power at the largest scales from $\ell = 2$ to $\ell \sim 7$. Fig. 5 shows a zoom on the TT panel of Fig. 4 for the lowest multipoles, where the estimates are now joined with broken lines in order to make this behavior clearer.




FIG. 3. Galactic masks. Upper panel: Masks we adopt in Temperature. Lower panel: Masks we adopt in Polarization. Blue region is for case “a”. Blue plus light blue region is for case “b”. Blue plus light blue and plus green region is for case “c”. Blue plus light blue plus green and plus orange region is for case “d”. See also Table II.

V. VARIANCE ANALYSIS

In order to measure such a deviation from WMAP 9 fiducial model we do not take into account the estimator given in Eq. (8) since it is not sensitive to a lack of power because of its quadratic expression in terms of distance from the theoretical model C_ℓ^{th} . Therefore we decide to naturally consider the expressions for the Variance of the CMB fluctuations in pixel space, which can be written as follows, see e.g. Appendix A of [31]

$$\langle \delta T^2 \rangle = \sum_{\ell \geq 2}^{\ell_{max}} \left(\frac{2\ell + 1}{4\pi} \right) C_\ell^{TT}, \quad (11)$$

$$\langle Q^2 \rangle = \sum_{\ell \geq 2}^{\ell_{max}} \left(\frac{2\ell + 1}{8\pi} \right) (C_\ell^{EE} + C_\ell^{BB}), \quad (12)$$

$$\langle U^2 \rangle = \langle Q^2 \rangle \quad (13)$$

where δT , is the fluctuation of the temperature map and Q and U are the Stokes parameters maps, see again e.g. [31] and references therein.




FIG. 4. WMAP 9 year CMB spectra at low resolution. Red symbols for case “a”, Black symbols for case “b”, Blue symbols for case “c” and Violet symbols for case “d”. See also Table II. From top to bottom, TT, EE, TE, BB, TB and EB spectrum. For each spectrum we show $\ell(\ell + 1)C_\ell/2\pi$ (given in μK^2) versus the multipole order ℓ . See also the text.




FIG. 5. Zoom of the TT spectrum given in Fig. 4 at the lowest multipoles. See also Table II.

A. Temperature analysis

In Fig. 6 we show the variance $\langle \delta T^2 \rangle$ computed through Eq. (11) with $\ell_{max} = 32$ for all the cases of Table II. Each panel of Fig. 6 shows the histogram obtained with MC simulations and the vertical bar corresponds to the WMAP data. Fig. 6 shows that increasing the Galactic mask, the WMAP TT variance is more and more anomalous. The probability to obtain a smaller value than WMAP from the random simulations is 9.87%, 2.2%, 0.4% and 0.01% for the case “a”, “b”, “c” and “d” respectively ⁶.

In Fig. 7 we plot such a percentage for all the cases of Table II as a function of ℓ_{max} . This Figure makes clear that the behavior given in Fig. 6 for $\ell_{max} = 32$ is in fact a general one and it holds for every choice of ℓ_{max} from 2 to 32. In particular, for the largest mask we consider, i.e. case “d”, the temperature variance is not consistent with Λ CDM model at 99.94% – 99.99% C.L. in the ℓ_{max} range [32, 40].

The low amplitude of the TT variance that we measure for the WMAP 9 year data, is dominated by the lowest multipoles. In order to show this, we plot in Fig. 8 the probability for the TT variance to obtain a smaller value than the WMAP one for the cases “a” (solid line), “b” (long dashed line), “c” (dashed line) and “d” (dotted line) of Table II as a function of ℓ_{min} with $\ell_{max} = 32$. Fig. 8 shows that excluding the quadrupole and the octupole in case “d” the anomaly disappears.

⁶ Note that the more interesting cases “a” and “d” are analyzed with 10^4 random simulations whereas “b” and “c” with 10^3 random simulations.




FIG. 6. TT Variance computed with Eq. (11) with $\ell_{max} = 32$ for the cases “a”, “b”, “c” and “d” from top to bottom and left to right respectively. See also Table II. Each panel shows counts versus $\langle \delta T^2 \rangle$. The vertical bars correspond to WMAP 9 year data. Upper left and lower right panels are built with 10^4 random extractions whereas upper right and lower left with 10^3 random extractions.




FIG. 7. The probability for TT variance to obtain a smaller value than the WMAP one for the cases “a” (solid line), “b” (long dashed line), “c” (dashed line) and “d” (dotted line) of Table II as a function of ℓ_{max} with $\ell_{min} = 2$.

B. Impact of the smoothing procedure in the temperature analysis

We check here the stability of our findings against the degradation procedure of the Temperature map, needed to perform the analysis at low resolution. Specifically we have degraded the WMAP 9 year ILC map to $N_{side} = 16$ performing no smoothing. Thanks to 10^4 MC simulations we have re-computed Eq. (11) for cases “a” and “d” with $\ell_{max} = 32$, see Fig. 9. The probability




FIG. 8. The probability for TT variance to obtain a smaller value than the WMAP one for the cases “a” (solid line), “b” (long dashed line), “c” (dashed line) and “d” (dotted line) of Table II as a function of ℓ_{min} with $\ell_{max} = 32$.




FIG. 9. TT Variance computed with Eq. (11) with $\ell_{max} = 32$ for the cases “a” and “d” from left to right respectively. See also Table II. Each panel shows counts versus $\langle \delta T^2 \rangle$. The vertical bars correspond to WMAP 9 year data of the ILC map pixellized at $N_{side} = 16$ without any smoothing. These panels are built with 10^4 random extractions.

to obtain a smaller value than WMAP from the random simulations is 5.02% and 0.01% for the case “a” and “d” respectively. Therefore our results are proven to be stable.

C. Polarization analysis

We have repeated for polarization the same analysis we have performed for temperature, see Eqs. (12),(13), finding no evident anomaly. For sake of brevity we report in Fig. 10 the analogous plot for polarization, already given in Fig. 7 for temperature. Even if in a regime that is noise dominated already at $\ell \gtrsim 10$, it is interesting to notice that increasing the polarization mask, the percentages move towards less anomalous values in a monotonic way with respect to the sky fraction (at least for $\ell_{max} \gtrsim 25$).




FIG. 10. The probability for $\langle Q^2 \rangle$ to obtain a smaller value than the WMAP one for the cases “a” (solid line), “b” (long dashed line), “c” (dashed line) and “d” (dotted line) of Table II as a function of ℓ_{max} .

VI. DISCUSSIONS AND CONCLUSIONS

Using a QML estimator, we have provided new APS estimates of WMAP 9 year data at large angular scales and studied their dependence on Galactic masking. We analyzed all six CMB spectra.

By using realistic MC simulations we have found that the lowest multipoles of the TT spectrum decrease in amplitude as the Galactic mask is increased. This indicates that the temperature anisotropies around the Galactic mask behave statistically differently from the anisotropies at high Galactic latitude. In particular, for the largest mask we consider, which covers $\sim 54\%$ of the sky, the temperature variance is not consistent with Λ CDM model at $99.94\% - 99.99\%$ C.L. in the ℓ_{max} range [32, 40].

We have also checked that the degradation procedure to obtain the low resolution WMAP 9 year ILC map is not impacting our findings. Specifically, we have degraded the WMAP 9 year ILC map to low resolution (corresponding to $N_{side} = 16$) performing no smoothing⁷ and we have re-evaluated for this map, Eq. (11) in cases “a” and “d”. Our results are also stable against the smoothing procedure, and therefore there is no evident spurious effect that is introduced in this data manipulation step. Furthermore we have verified that the lowest multipoles are responsible of such a variance anomaly.

Though based on different estimators, our findings are in agreement with the analysis performed in [3], for WMAP 5 and 7 data, and with [18], where WMAP 3 year data is considered.

⁷ We remind that our results presented in Subsection V A are based on the WMAP 9 year ILC map smoothed with FWHM=9.1285 degrees, see Section III.

We do not find evidence of a similar effect in the other polarized spectra. This is not surprising since they are noise dominated already at $\ell \sim 10$.

Recently PLANCK analysis of the variance for the temperature CMB anisotropies has appeared [26]. We aim at analyzing these data as well along the same lines as given here for the temperature case since the pattern of TT APS at large angular scales is close to the WMAP 9 year data when the same mask is adopted [22]. Note that the lack of power at large angular scales claimed in [22] is not the same as the one discussed here, since it involves different range of multipoles (see Fig. 39 of [22]).

In conclusion, we speculate that this “low temperature variance anomaly” might be an indication of a non-standard phase of the cosmological evolution of the Inflaton before the slow roll phase as studied for example in [4], [6] and [5] for different pre-inflationary scenarios⁸.

ACKNOWLEDGMENTS

A.G. wishes to thank Belen Barreiro, Marcos Cruz, Enrique Martinez-Gonzales and Patricio Vielva for fruitful conversations. We acknowledge the use of computing facilities at NERSC (USA) and CINECA (ITALY). We acknowledge use of the HEALPix (Gorski et al. 2005) software and analysis package for deriving the results in this paper. We acknowledge the use of the Legacy Archive for Microwave Background Data Analysis (LAMBDA). Support for LAMBDA is provided by the NASA Office of Space Science. Work supported by ASI through ASI/INAF Agreement I/072/09/0 for the Planck LFI Activity of Phase E2 and by MIUR through PRIN 2009 (grant n. 2009XZ54H2).

-
- [1] C. L. Bennett, D. Larson, J. L. Weiland, N. Jarosik, G. Hinshaw, N. Odegard, K. M. Smith and R. S. Hill *et al.*, arXiv:1212.5225 [astro-ph.CO].
 - [2] P. Bielewicz, A. J. Banday and K. M. Gorski, arXiv:1111.6046 [astro-ph.CO].
 - [3] M. Cruz, P. Vielva, E. Martinez-Gonzalez and R. B. Barreiro, arXiv:1005.1264 [astro-ph.CO].
 - [4] C. Destri, H. J. de Vega and N. G. Sanchez, Phys. Rev. D **81**, 063520 (2010) [arXiv:0912.2994 [astro-ph.CO]].
 - [5] S. Downes and B. Dutta, arXiv:1211.1707 [hep-th].
 - [6] E. Dudas, N. Kitazawa, S. P. Patil and A. Sagnotti, JCAP **1205**, 012 (2012) [arXiv:1202.6630 [hep-th]].

⁸ In fact models of compact Universes lead to similar effect in the Temperature maps but they also predict the existence of specific patterns like “circles-in-the-sky” that have not been detected, see e.g. [2].

- [7] J. Dunkley *et al.* [WMAP Collaboration], *Astrophys. J. Suppl.* **180**, 306-329 (2009). [arXiv:0803.0586 [astro-ph]].
- [8] K.M. Gorski, E. Hivon, A.J. Banday, B.D. Wandelt, F.K. Hansen, M. Reinecke and M. Bartelmann, 2005, HEALPix: A Framework for High-resolution Discretization and Fast Analysis of Data Distributed on the Sphere, *Ap.J.*, 622, 759-771
- [9] J. Grain, M. Tristram and R. Stompor, arXiv:0903.2350 [astro-ph.CO].
- [10] A. Gruppuso, A. De Rosa, P. Cabella, F. Paci, F. Finelli, P. Natoli, G. de Gasperis, N. Mandolesi, *Mon. Not. Roy. Astron. Soc.* **400**, 463-469 (2009). [arXiv:0904.0789 [astro-ph.CO]].
- [11] A. Gruppuso, F. Finelli, P. Natoli, F. Paci, P. Cabella, A. De Rosa and N. Mandolesi, *Mon. Not. Roy. Astron. Soc.* **411**, 1445 (2011) [arXiv:1006.1979 [astro-ph.CO]].
- [12] A. Gruppuso, P. Natoli, N. Mandolesi, A. De Rosa, F. Finelli and F. Paci, *JCAP* **1202**, 023 (2012) [arXiv:1107.5548 [astro-ph.CO]].
- [13] G. Hinshaw, D. Larson, E. Komatsu, D. N. Spergel, C. L. Bennett, J. Dunkley, M. R. Nolta and M. Halpern *et al.*, “Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Parameter Results,” arXiv:1212.5226 [astro-ph.CO].
- [14] E. Hivon *et al.*, *Astrophys. J.* **567**, 2 (2002)
- [15] N. Jarosik *et al.* [WMAP Collaboration], *Astrophys. J. Suppl.* **170**, 263 (2007) [astro-ph/0603452].
- [16] D. Larson, J. Dunkley, G. Hinshaw, E. Komatsu, M. R. Nolta, C. L. Bennett, B. Gold, M. Halpern *et al.*, *Astrophys. J. Suppl.* **192**, 16 (2011). [arXiv:1001.4635 [astro-ph.CO]].
- [17] D. Molinari *et al.* (2013), to appear.
- [18] C. Monteserin, R. B. B. Barreiro, P. Vielva, E. Martinez-Gonzalez, M. P. Hobson and A. N. Lasenby, *Mon. Not. Roy. Astron. Soc.* **387**, 209 (2008) [arXiv:0706.4289 [astro-ph]].
- [19] F. Paci, A. Gruppuso, F. Finelli, P. Cabella, A. De Rosa, N. Mandolesi and P. Natoli, *Mon. Not. Roy. Astron. Soc.* **407**, 399 (2010) [arXiv:1002.4745 [astro-ph.CO]].
- [20] F. Paci, A. Gruppuso, F. Finelli, A. De Rosa, N. Mandolesi and P. Natoli, arXiv:1301.5195 [astro-ph.CO].
- [21] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1303.5062 [astro-ph.CO].
- [22] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1303.5075 [astro-ph.CO].
- [23] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1303.5076 [astro-ph.CO].
- [24] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1303.5080 [astro-ph.CO].
- [25] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1303.5082 [astro-ph.CO].
- [26] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1303.5083 [astro-ph.CO].
- [27] G. Polenta *et al.*, *JCAP* **0511**, 001 (2005) [arXiv:astro-ph/0402428].
- [28] R. Saha, P. Jain and T. Souradeep, *Astrophys. J.* **645**, L89 (2006) [arXiv:astro-ph/0508383].
- [29] F. Schiavon, F. Finelli, A. Gruppuso, A. Marcos-Caballero, P. Vielva, R. G. Crittenden, R. B. Barreiro and E. Martinez-Gonzalez, arXiv:1203.3277 [astro-ph.CO].
- [30] M. Tegmark, *Phys. Rev. D* **55**, 5895 (1997) [astro-ph/9611174].

[31] M. Tegmark and A. de Oliveira-Costa, Phys. Rev. D **64** (2001) 063001 [astro-ph/0012120].